

- Baaritaanka jirka ee ee cudurka qaaxada waxaa loo sameeyaa si loo ogaado jeermiga cudurka qaaxada ee jirkaada ku jiro.
- Baarista jirka kuma ogaan kartid haddii uu ku hayo cudurka qaaxada.
- Haddii baaritaanka jirka (maqaarka) laga helo cudurka qaaxada ama tiibishada waxaad u baahan doontaabaaritaanno kale.




<http://www.vdh.state.va.us/epi/tb>

[“What is a TB Skin Test?”]
Somali

November 8, 2002

Waa maxay
baarista
TB-da ee
jirka
(maqaarka)
?





Baarista qaaxada (TB- da) waa baaritaanka jirka (maqaarka) oo kaliya!

Sidee baarista loo
sameeyaa?

Waxaa la isticmaalaa cirbad yar oo
lagu duro maqaarka gacanta, oo ay
ku jirto walax baarista la yiraahdo
tuberculin. TB-da kama qaadi
kartid walaxda baaristan.

Waa in aad ku soo noqotaaa
muddo 2 ilaa 3 maalmood ah, si
shaqaala caafimaadku uu u fiiriyo
gacaantaada.



Maxay natijadaydu
macnaheedu tahay?

Haddii ay gacanta ku soo yar
bararto, waxaa laga yaabaa in
baaristu tahay hubaal “boosatifo”.
Shaqaalahaa caafimaadku ayaa
go’aminaya in cudurka qaaxada
aad qabto marka ay cabiraan inta
bararku la egyptay. Haddii

baarista ay cadeyso in cudurka aad
qabto oo “boosatifo” aad tahay,
waxaad u baahantahay baaris
dheeraad ah, sida raajada feeraha ,
si loo ogaado in uu ku hayo
cudurka qaaxada ama aad la
bugto.



Hadii natijadu aanay hubaal
ahayn “negatiifo”, oo gacantu
aysan ku soo bararin, cudurka
TB-da ma qabtid. Tani
macnaheedu waxay tahay in
laga yaabo in jirkaada uusan ku
jirin jeermiska TBda.

LAAKIIN, laga yaabee inaad
u baahan tahay baaritaan kale,
haddii aad ka ag dhawayd qof
la jiran cudurka qaaxada ama
tiibishada).